

SOT Therapy – Virus/Lyme ONLY

Pre and Post Administration Guidelines

The purpose of these guidelines is to ensure the highest level of effectiveness of the SOT therapy by removing treatments that interfere with and/or diminish the effectiveness of that therapy.

**** Adherence to these guidelines will improve therapy effectiveness and patient outcomes.**

For Virus/Lyme (V. Antagonist) SOT:

- **Pre-SOT Administration:** The patient must be off ALL antiviral/antibacterial/ antiparasitic medications and therapies for **14 days** prior to the administration of the therapy
 - Antibiotics (oral or IV)
 - Antiviral medications (oral or IV)
 - Anti-parasitic medications (oral or IV)
 - Vitamin C (IV)
 - Colloidal Silver (oral or IV)
 - H2O2 IV
 - Ozone (IV)
 - Natural Substances (ALL oral supplements that are trying to kill or suppress the same target as the SOT)
- **Day of SOT Administration:** Absolutely NO other therapies are recommended on the day of the SOT administration.
- **Post SOT Administration:** All therapies can be resumed **7 days** after the administration to allow the SOT time to locate its target without interference.

Note: Detox supplements such as binders, glutathione, cytokine reducers, etc. will not interfere with the action of the SOT and can be used both pre and post SOT administration.